

Grain Acceleration by MHD Turbulence: Gyroresonance Mechanism

Huirong Yan and A. Lazarian

Department of Astronomy, University of Wisconsin, 475 N. Charter St., Madison, WI 53706; yan, lazarian@astro.wisc.edu

ABSTRACT

We discuss a new mechanism of dust acceleration that acts in a turbulent magnetized medium. The magnetohydrodynamic (MHD) turbulence includes both fluid motions and magnetic fluctuations. We show that while the fluid motions bring about grain motions through the drag, the electromagnetic fluctuations, can accelerate grains through resonant interactions. In this paper we calculate grain acceleration by gyroresonance in Cold Neutral Medium. We consider both incompressible and compressible MHD modes. We show that fast modes dominate grain acceleration. For the parameters chosen, fast modes render supersonic velocities to grains that may shatter grains and enable efficient absorption of heavy elements. Since the grains are preferentially accelerated with large pitch angles, the supersonic grains get aligned with long axes perpendicular to the magnetic field.

Subject headings: dust, extinction–ISM: particle acceleration–kinematics and dynamics–magnetic fields

1. INTRODUCTION

Dust is an important constituent that is essential for heating and cooling of the interstellar medium (ISM). It interferes with observations in the optical range, but provides an insight to star-formation activity through far-infrared radiation. It also enables molecular hydrogen formation and traces the magnetic field via emission and extinction polarimetry (see Lazarian 2003). The basic properties of dust (optical, alignment etc.) strongly depend on its size distribution. The latter evolves as the result of grain collisions, whose frequency and consequences (coagulation, cratering, shattering, and vaporization) depend on grain relative velocities (see discussions in Draine 1985, Lazarian & Yan 2002a,b).

It is known that all these problems require our understanding of grain motions in turbulent interstellar medium. Although turbulence has been invoked by a number of authors (see Kusaka et al. 1970, Draine 1985, Ossenkopf 1993, Weidenschilling & Ruzmaikina 1994) to provide substantial grain relative motions, the turbulence they discussed was not magnetized. In a recent paper (Lazarian & Yan 2002a, hereafter LY02) we applied the theory of Alfvénic turbulence (Goldreich & Sridhar 1995, henceforth GS95, Cho, Lazarian & Vishniac 2002a for a review) to grain acceleration through gaseous drag.

Here we account for the acceleration that arises from the resonant interaction of charged grains with MHD turbulence. To describe the turbulence statistics we use the analytical fits to the statistics of Alfvénic modes obtained in Cho, Lazarian & Vishniac (2002b, hereafter CLV02) and compressible modes obtained in Cho & Lazarian (2002, hereafter CL02).

2. Acceleration of Grains by Gyroresonance

Turbulent acceleration may be viewed as the acceleration by a spectrum of MHD waves that can be decomposed into incompressible Alfvénic, and compressible fast and slow modes (see CL02). There exists an important analogy between dynamics of charged grains and dynamics of cosmic rays (see Yan & Lazarian 2002, henceforth YL02) and we shall modify the existing machinery used for cosmic rays to describe charged grain dynamics¹. The energy exchange involves resonant interactions between the particles and the waves. Specifically, the resonance condition is $\omega - k_{\parallel}v\mu = n\Omega$, ($n = 0, \pm 1, \pm 2, \dots$), where ω is the wave frequency, $k_{\parallel}$ is the parallel component of wave vector $\mathbf{k}$ along the magnetic field, v is the particle velocity, μ is the cosine of the pitch angle relative to the magnetic field, $\Omega = qB/(mc)$ is the Larmor frequency of the particle. The sign of n denotes the polarization of the wave. $+$ represents left hand polarization and $-$ is for right hand polarization. Basically there are two main types of resonant interactions: gyroresonance acceleration and transit acceleration. Transit acceleration ($n = 0$) requires longitudinal motions and only operates with compressible modes. It happens when $k_{\parallel}v\mu = \omega$. As the phase speed is $V_{fast} \geq V_A$ for fast waves, where V_A is the Alfvén speed, it is clear that it can be only applicable to super-Alfvénic grains that we do not deal with here.

Gyroresonance occurs when the Doppler-shifted frequency of the wave in the grain’s guiding center rest frame $\omega_{gc} = \omega - k_{\parallel}v\mu$ is a multiple of the particle gyrofrequency, and the rotating direction of wave electric vector is the same as the direction for Larmor gyration

¹In what follows we assume that the time scale for grain charging is much shorter than the grain Larmor period.

of the grain. The gyroresonance scatters and accelerates the particles. The efficiency of the two processes for charged grains can be described by the Fokker-Planck coefficients $D_{\mu\mu}$ and D_{pp}/p^2 , where p is the particle momentum (see Schlickeiser & Achatz 1993, YL02). The ratio of the two rates depends on the ratio of the particle velocity, the Alfvén speed and pitch angle, $p^2 D_{\mu\mu}/D_{pp} = [(v\zeta/V_A) + \mu]^2$, where $\zeta = 1$ for Alfvén waves and $\zeta = k_{\parallel}/k$ for fast modes. We see that the scattering is less efficient for sub-Alfvénic grains unless most particles move parallel to the magnetic field. We shall show later that as the result of acceleration, μ will tend to 0. Therefore in the zeroth order approximation, we ignore the effect of scattering and assume that the pitch angle cosine μ does not change while being accelerated. In this case, the Fokker-Planck equation, which describes the diffusion of grains in the momentum space, can be simplified (see Pryadko & Petrosian 1997):

$$\frac{\partial f}{\partial t} + v\mu \frac{\partial f}{\partial z} = \frac{1}{p^2} \frac{\partial}{\partial p} p^2 D_{pp}(\mu) \frac{\partial f}{\partial p}, \quad (1)$$

where f is the distribution function. Apart from acceleration, a grain is subjected to gaseous friction. For the sake of simplicity we assume that grains are moving in respect to stationary gas and turbulence provides nothing but the electromagnetic fluctuations. The acceleration of grains arising from the gas motion is considered separately (see LY02). We describe the stochastic acceleration by the Brownian motion equation: $mdv/dt = -v/S + Y$, where m is the grain mass, Y is the stochastic acceleration force, $S = t_{drag}/m$ is the mobility coefficient, t_{drag} is the gas drag time (Draine 1985, LY02). The drag time due to collisions with atoms is essentially the time for collisions with the amount of gas with the mass equal to that of the grain, $t_{drag}^0 = (a\rho_{gr}/n_n)(\pi/8m_n k_B T)^{1/2}$, where a is the grain size, T is the temperature, ρ_{gr} is the mass density of grain. We adopt $\rho_{gr} = 2.6\text{gcm}^{-3}$ for silicate grains. The ion-grain cross-section due to long-range Coulomb force is larger than the atom-grain cross-section. Therefore, in the presence of collisions with ions, the effective drag time decreases by the factor given in Draine & Salpeter (1979). When the grain velocity gets supersonic (Purcell 1969), the gas drag time is given by $t_{drag}^s = t_{drag}/(0.75 + 0.75c_s^2/v^2 - c_s^3/2v^3 + c_s^4/5v^4)$, where c_s is the sound speed.

Multiply the Brownian equation above by v and take the ensemble average, we obtain

$$m \frac{d \langle v^2 \rangle}{dt} = - \frac{\langle v^2 \rangle}{S} + \langle \dot{\epsilon} \rangle, \quad (2)$$

Following the approach similar to that in Melrose (1980), we can get from Eq.(1) the energy gain rate $\langle \dot{\epsilon} \rangle$ for the grain with pitch angle μ

$$\langle \dot{\epsilon} \rangle = \frac{1}{p^2} \frac{\partial}{\partial p} (vp^2 D_{pp}(\mu)), \quad (3)$$

where the Fokker-Planck coefficient $D_{pp}(\mu)$ is given below.

Adopting the result from quasi-linear theory (hereafter QLT, see Schlickeiser & Achatz 1993), the momentum diffusion coefficient is (see YL02)

$$\begin{aligned} D_{pp}(\mu) = & \frac{\pi\Omega^2(1-\mu^2)p^2V_A^2}{2v^2} \mathcal{R} \int dk^3 \frac{\tau_k^{-1}}{\tau_k^{-2} + (\omega - k_{\parallel}v\mu - n\Omega)^2} \\ & [(J_{n+1}^2(\frac{k_{\perp}v_{\perp}}{|\Omega|})K_{\mathcal{R}\mathcal{R}}(\mathbf{k})) + J_{n-1}^2(\frac{k_{\perp}v_{\perp}}{|\Omega|})K_{\mathcal{L}\mathcal{L}}(\mathbf{k})) \\ & - J_{n+1}(\frac{k_{\perp}v_{\perp}}{|\Omega|})J_{n-1}(\frac{k_{\perp}v_{\perp}}{|\Omega|}) \\ & (e^{i2\phi}K_{\mathcal{R}\mathcal{L}}(\mathbf{k}) + e^{-i2\phi}K_{\mathcal{L}\mathcal{R}}(\mathbf{k}))], \end{aligned} \quad (4)$$

where τ_k is the nonlinear decorrelation time and essentially the cascading time of the turbulence, $k_{\perp} = \sqrt{k_x^2 + k_y^2}$ is the perpendicular component of the wave vector, $v_{\perp}$ is the perpendicular component of grain velocity, $\phi = \tan^{-1}(k_y/k_x)$, $K_{\alpha\beta}(\mathbf{k})$ is the velocity correlation tensor and will be given in §3, $\mathcal{L}, \mathcal{R} = (x \pm iy)/\sqrt{2}$ represent left and right hand polarization. At the magnetostatic limit, $\tau_k \rightarrow \infty$, the so-called Breit-Wigner-type function transfers into δ function, i.e., $\tau_k^{-1}/(\tau_k^{-2} + (\omega - k_{\parallel}v\mu - n\Omega)^2) \rightarrow \pi\delta(\omega - k_{\parallel}v\mu - n\Omega)$. Magnetostatic limit is correct for fast moving particles (see YL02), but for sub-Alfvénic grains we should use Eq.(4). However, we should not integrate over the whole range of $k_{\parallel}$, because the contribution from large scale is spurious (see YL02). This contribution stems from the fact that in QLT, an unperturbed particle orbit is assumed, which results in non-conservation of the adiabatic invariant $\xi = mv_{\perp}^2/2B_0$, where B_0 is the large-scale magnetic field. Noticing that the adiabatic invariant is conserved only when the electromagnetic field varies on a time scale larger than $|\Omega|^{-1}$, we truncate our integral range, namely, integrate from k_{res} instead of the injection scale L^{-1} . For Alfvénic turbulence $\omega = |k_{\parallel}|V_A$, the resonant scale corresponds to $|k_{\parallel, res}| = |\Omega/(V_A - v\mu\zeta)|$. For fast modes in a low β medium (where $\beta \equiv P_{gas}/P_{mag} = 2c_s^2/V_A^2$ is the ratio of gaseous pressure and magnetic pressure), $\omega = kV_A$, the resonant scale is $k_{res} = I\Omega/(V_A - v\mu\zeta)$. The upper limit of the integral k_c is set by the dissipation of the MHD turbulence, which varies with the medium.

Integrating from k_{res} to k_c , we can obtain from Eq.(4) and (3) the energy gain rate $\langle \dot{\epsilon} \rangle$ as a function of v and μ . Then with the $\langle \dot{\epsilon} \rangle$ known, we can estimate the grain acceleration. Solving the Eq.(2) iteratively, we can get the grain velocity as a function of time. We check that the grain velocities converge to a constant value after the drag time.

Thus inserting the acceleration rate by fast and Alfvén modes into Eq.(2), we can obtain the final grain velocities as a function of μ . As $\langle \epsilon \rangle$ increases with pitch angle, grains gain the maximum velocities perpendicular to the magnetic field and therefore the averaged μ decreases.

3. MHD Turbulence and Its Tensor Description

Unlike hydrodynamic turbulence, Alfvénic turbulence is anisotropic, with eddies elongated along the magnetic field. The Alfvénic turbulence is described by GS95 model which postulates that $k_{\perp} v_k \sim k_{\parallel} V_A$. This may be viewed as coupling of eddies perpendicular to the magnetic field and wave-like motions parallel to the magnetic field. For magnetically dominated, the so-called low β plasma, CL02 showed that the coupling of Alfvén and compressible modes is weak and that the Alfvén and slow modes follow the GS95 spectrum. This is consistent with the analysis of HI velocity statistics (Lazarian & Pogosyan 2000, Stanimirovic & Lazarian 2001). According to CL02, fast modes are isotropic. In what follows, we consider both Alfvén modes and compressible modes in low β plasma.

Within the random-phase approximation, the velocity correlation tensor in Fourier space is (see Schlickeiser & Achatz 1993) $\langle v_{\alpha}(\mathbf{k}, t) v_{\beta}^*(\mathbf{k}', t + \tau) \rangle / V_A^2 = \delta(\mathbf{k} - \mathbf{k}') K_{\alpha\beta}(\mathbf{k}) e^{-\tau/\tau_k}$, where $v_{\alpha,\beta}$ is the time-dependent velocity fluctuation in $\mathbf{k}$ space associated with the turbulence.

The velocity correlation tensor for Alfvénic turbulence is (CLV02),

$$\begin{aligned} K_{ij}(\mathbf{k}) &= \frac{L^{-1/3}}{12\pi} I_{ij} k_{\perp}^{-10/3} \exp(-L^{1/3} |k_{\parallel}| / k_{\perp}^{2/3}), \\ \tau_k &= (L/V_A)(k_{\perp} L)^{-2/3} \sim (k_{\parallel} V_A)^{-1} \end{aligned} \quad (5)$$

where $I_{ij} = \{\delta_{ij} - k_i k_j / k^2\}$ is a 2D tensor in the $x - y$ plane which is perpendicular to the magnetic field, L is the injection scale, V is the velocity at the injection scale. Velocity fluctuations related to slow modes are subdominant for magnetically dominated plasmas (CL02) and we do not consider them.

Fast modes are isotropic and have one dimensional energy spectrum $E(k) \propto k^{-3/2}$ (CL02). In low β medium, the velocity fluctuations are always perpendicular to $\mathbf{B}_0$ for all $\mathbf{k}$, the corresponding correlation is (YL02)

$$K_{ij}(\mathbf{k}) = \frac{L^{-1/2}}{8\pi} J_{ij} k^{-7/2}, \quad \tau_k = (k/L)^{-1/2} \times V_A/V^2 \quad (6)$$

where $J_{ij} = k_i k_j / k_\perp^2$ is also a 2D tensor in $x - y$ plane.

4. Results

We consider a typical cold neutral medium (CNM), $T = 100\text{K}$, $n_n = 30\text{cm}^{-3}$, $B_0 = 6.3\mu\text{G}$. Here, we only consider large grains ($10^{-6}\text{cm} < a < 10^{-4}\text{cm}$), which carry most grains mass ($\sim 80\%$) in ISM. The mean grain charge was obtained from the average electrostatic potential $\langle U \rangle$ in Weingartner & Draine (2001). MHD turbulence requires that fluid velocities are smaller than the Alfvén speed. Therefore we assume that the injection of energy happens at the scale L where the equipartition between magnetic and kinetic energies, i.e., $V = V_A$, is reached. We assume that the velocity dispersion at the scale $l = 10\text{pc}$ is 5km/s and that the turbulence at large scales proceeds in tenuous warm media with Alfvén speed larger or equal to 5km/s . In partially ionized medium, a viscosity caused by neutrals results in decoupling on the characteristic time scale (see LY02)

$$t_{damp} \sim \nu_n^{-1} k^{-2} \sim (l_n v_n)^{-1} k^{-2}, \quad (7)$$

where ν_n is the kinetic viscosity, l_n is the neutral mean free path, v_n is the thermal velocity of neutrals. Given the parameters above, $l_n = 7 \times 10^{12} \text{ cm}$. When its cascading rate $\tau_k^{-1} = k_\parallel V_A \sim k_\perp^{2/3} L^{-1/3} V_A$ (see Eq.5) equals to the damping rate² $t_{damp}^{-1}(k_\perp)$, Alfvénic turbulence is assumed damped³. This defines the cutoff wave number of the turbulence $k_{\parallel,c} = 2.4 \times 10^{-16} \text{ cm}^{-1}$ and the time scale $\tau_c = (k_{\parallel,c} V_A)^{-1} = 1.7 \times 10^{10} \text{ s}$. Assuming that the grain velocities are smaller than Alfvén speed, we can find that the prerequisite for the gyroresonance $|k_{\parallel,c}| > |k_{\parallel,res}|$ is the same as $\tau_L = 2\pi/|\Omega| > \tau_c$, the condition for effective hydro drag (see LY02). Thus we see that Alfvén modes cannot accelerate grains (with $a < 2 \times 10^{-5} \text{ cm}$) through gyroresonance unless the velocities of these grains are already super-Alfvénic. The cutoff of fast modes corresponds to the scale where the cascading time scale $\tau_k = t_{damp}$ and this gives the cutoff wave number $k_c = 4.9 \times 10^{-15} \text{ cm}^{-1}$. In the present paper we consider neutral gas of low ionization, and therefore the damping owing to ions, including collisionless damping, is disregarded (compare to YL02). Using the procedure described in §2, we obtain the grain velocities for different pitch angles. The acceleration is maximal in the direction perpendicular to the magnetic field. Those values are presented in

²As pointed out in §2, the eddy motions happen in the direction perpendicular to the magnetic field, so $k_\perp$ is used to calculate t_{damp} for Alfvén modes.

³Thus we ignore the effect of slowly evolving magnetic structures associated with a recent reported new regime of turbulence below the viscous damping cutoff (Cho, Lazarian & Vishniac 2002c).

Fig.1. If averaged over μ , the velocities are smaller by less than 20%.

In order to compare different processes, we account for the hydro drag (see LY02 for a discussion of Alfvén-induced drag). The fast modes cause the relative movement of the grain to the ambient gas also by gaseous drag. Unlike gyroresonance, this relative motion arises from the decoupling from the gas. At large scale grains are coupled with the ambient gas, and the slowing fluctuating gas motions will only cause an overall advection of the grains with the gas (Draine 1985), which we are not interested. The largest velocity difference occurs on the largest scale where grains are still decoupled. While in the hydrodynamic case the decoupling happens on the time scale t_{drag} , the grains are constrained by the Larmor gyration in MHD case unless $t_{drag} < \tau_L$ (see LY02). The latter condition is true for high density gas, however (see Lazarian & Yan 2002b). The velocity fluctuations for fast modes scales as $v_k \propto k^{-1/4} \propto \omega^{-1/4}$, where $\omega = kV_A$ is the frequency of fast modes (see Eq.(6)). In low β medium, the velocity fluctuations are perpendicular to the magnetic field. Therefore, grains get velocity dispersions perpendicular to the magnetic field, $v \simeq V(\tau_L/\tau_{max})^{1/4}$, where $\tau_{max} = L/V$ is the time scale at the injection scale of turbulence. Then we also need to consider the effect of damping. Similar to Alfvén modes, the condition for effective hydro drag $\tau_L > \tau_c = 2\pi/\omega_c = 5 \times 10^9$ s is the same as $k_c > k_{res}$ for sub-Alfvénic particles, which is the requirement for gyroresonance. Our calculation shows that the corresponding grain size is 4×10^{-6} cm. For smaller grains, their velocities are reduced $v \sim v_c \times \tau_L/\tau_c \sim V(\tau_L/\tau_{max})^{1/4}(\tau_L/\tau_c)^{3/4}$, where v_c is the velocity of turbulence at the damping scale (see Lazarian & Yan 2002b). In Fig.1 we plot the velocity of grain with $\mu = 0$ as a function of grain size since all the mechanisms preferentially accelerate grains in this direction.

How would the results vary as the parameters of the partially ionized medium vary? From Eqs.(2), (3)&(4), we find the grain velocity is approximately equal to $v \sim (< \dot{\epsilon} > t_{drag}/m)^{1/2} \sim 2.5 \times 10^4 L_{10}^{-1/4} V_5 B_\mu^{1/2} (q_e/a_5^3)^{0.3} t_{drag}^{0.5}$ cm/s, where $L_{10} = L/10$ pc is the injection scale defined above, $V_5 = V_A/5$ km s $^{-1}$, $B_\mu = B/1\mu$ G, $q_e = q/1$ electron. Noticing that the hydro drag by fast modes decreases with the magnetic field, $v \propto B^{-1/4}$ beyond the damping cutoff and $v \propto B^{-1}$ below the cutoff (see the last paragraph), we see that the relative importance of the gyroresonance and hydro drag depends on the magnitude of the magnetic field.

It has been shown that the composition of the galactic cosmic ray seems to be better correlated with volatility of elements (Ellison, Drury & Meyer 1997). The more refractory elements are systematically overabundant relative to the more volatile ones. This suggests that the material locked in grains must be accelerated more efficiently than gas-phase ions (Epstein 1980; Ellison, Drury & Meyer 1997). The stochastic acceleration of grains, in this case, can act as a preacceleration mechanism.




Fig. 1.— Grain velocity vs. size owing to different acceleration processes in CNM. The solid line represents the gyroresonance with fast modes. The dashdot line refers to the gyroresonance with Alfvén modes. Gyroresonance with both modes works only for large grains owing to the cutoff by viscous damping. The cutoff scales for fast and Alfvén modes are different mostly due to the anisotropy of Alfvén modes. The dotted line is the result from hydro drag with Alfvén modes (see LY02), the dashed line represents the hydro drag with fast modes. The corresponding grain Larmor frequency Ω and t_{drag}^{-1} are also given along the abscissas.

Grains moving supersonically can also efficiently vacuum-clean heavy elements as suggested by observations (Wakker & Mathis 2000). Grains can also be aligned if the grains get supersonic (see review by Lazarian 2003). Indeed, the scattering is not efficient for slowly moving grains so that we may ignore the effect of scattering to the angular distribution of the grains. Since the acceleration of grains increases with the pitch angle of the grain (see Eqs.(3) and (4)), the supersonic grain motions will result in grain alignment with long axes perpendicular to the magnetic field.

It is believed that silicate grains won't be shattered unless their velocities reach 2.7km/s (Jones et al. 1996). Nevertheless, the threshold velocities depend on the velocity of turbulence at the injection scale and on the grain structure, i.e., solid or fluffy. Thus shattering of the largest grains is possible.

5. SUMMARY

In the paper we showed that

1. Fast modes provide the dominant contribution for the acceleration of charged grains. The velocities obtained are sufficiently high to be important for shattering large grains and efficiently absorbing heavy elements from gas. Alfvén modes are not important because of their anisotropy.
2. Depending on the relative importance of the magnetic field, gyroresonance (strong B) or hydro drag (weak B) by fast modes dominates grain acceleration. For small grains, hydro drag by fast modes is the most important while the gyroresonance is not present as the turbulence at the resonant frequencies gets viscously damped.
3. In low β medium, all the mechanisms tend to preferentially accelerate grains in the direction perpendicular to the magnetic field. Among them, gyroresonance with fast modes can render grains with supersonic motions, which can result in grain alignment perpendicular to the magnetic field.

We acknowledge stimulating communications with E. Zweibel, which were very important at the initial stage of this work. We thank J. Mathis, J. Cho and R. Schlickeiser for helpful discussions. This work is supported by NSF grant AST01-25544.

REFERENCES

- Cho, J. & Lazarian, A. 2002, Phys. Rev. Lett., 88, 245001 (CL02)
- Cho, J., Lazarian, A. & Vishniac, E. T. 2002a, in Turbulence and Magnetic field in Astrophysics, Eds. T. Passot & E. Falgarone (Springer LNP), p.56
- Cho, J., Lazarian, A. & Vishniac, E. T. 2002b, ApJ, 564, 291 (CLV02)
- Cho, J., Lazarian, A. & Vishniac, E. T. 2002c, ApJ, 566, 49L
- Draine, B. T. 1985, in Protostars and Planets II, ed. D. C. Black & M. S. Matthews (Tucson: Univ. Arizona Press), p.621
- Draine, B. T. & Salpeter, E. E. 1979, ApJ, 231, 77
- Ellison, D. C., Drury, L. O’C. & Meyer, J.-P. 1997, ApJ, 487, 197
- Epstein, R. I. 1980, MNRAS, 193, 723
- Goldreich, P. & Sridhar, H. 1995, ApJ, 438, 763 (GS95)
- Jones, A. P., Tielens, A. G. G. M. & Hollenbach, D. J. 1996, ApJ, 469, 740
- Kusaka, T., Nakano, T., & Hayashi, C., 1970, Prog. Theor. Phys., 44, 1580
- Lazarian, A. 2003, J. Quant. Spectrosc. Radiat. Transfer, 79, 881
- Lazarian, A. & Pogosyan, D. 2000, ApJ, 537, 720
- Lazarian, A. & Yan, H. 2002a, ApJ, 566, 105L (LY02)
- Lazarian, A. & Yan, H. 2002b, astro-ph/0205283
- Mathis, J. S. 1990, ARA&A, 28, 37
- Melrose, D. B. 1980, Plasma Astrophysics (NY: Gordon & Breach).
- Ossenkopf, V. 1993, A&A 280, 617
- Pryadko, J. M. & Petrosian, V. 1997, ApJ, 482, 774
- Purcell, E. M. 1969, Physica 41, 100
- Schlickeiser, R. & Achatz, U. 1993, J. Plasma Phy. 49, 63
- Stanimirovic, S. & Lazarian, A. 2001, ApJ, 551, L53
- Wakker, B. P. & Mathis, J. S. 2000, ApJ, 544, 107L
- Weidenschilling, S. J. & Ruzmaikina, T. V. 1994, ApJ, 430, 713
- Weingartner, J. C. & Draine, B. T. 2001, ApJs, 134, 263
- Yan, H. & Lazarian, A. 2002, Phy. Rev. Lett, 89, 281102 (YL02)